Examiner. Withdrawal of all rejections and reconsideration of the amended claims is requested. An early allowance is earnestly sought.

Respectfully submitted,

Dated: July 3, 2003

Rochelle K. Seide Patent Office Reg. No. 32,300

Carmella L. Stephens Patent Office Reg. No. 41,328

BAKER BOTTS L.L.P. 30 Rockefeller Plaza New York, New York 10112-0228

Attorneys for Applicants (212) 408-2500

APPENDIX

IN THE CLAIMS:

Please cancel claims 1-7, 12, 14 and 20 and insert the following new claims:

21. (new) A method for generating a hepatic cell culture comprising:

co-culturing hepatocytes and nonparenchymal cells derived from disaggregated liver tissue, in the presence of

(a) one or more growth factors that support the growth of hepatocytes and

(b) a matrix wherein said matrix is a bead coated with at least one
biologically active molecule that promotes cell adhesion
under conditions sufficient to allow for the proliferation of said hepatocytes while
retaining hepatic function of said hepatocytes.

22. (new) A method for generating a hepatic cell culture comprising:

established hepatic cell lines comprising hepatocytes and non-

parenchymal cells, in the presence of

(a) one or more growth factors that support the growth of

hepatocytes and

(b) a matrix wherein said matrix is a bead coated with at least one

biologically active molecule that promotes cell adhesion

under conditions sufficient to allow for the proliferation of said hepatocytes while retaining hepatic function of said hepatocytes.

23. (new) The method of claim 21 or 22 wherein the matrix is in the form of
polystyrene beads.
24. (new) The method of claim 21 or 22 wherein the matrix is coated with an
extracelluar matrix protein.
25. (new) The method of claim 21 or 22 wherein the matrix is coated with type I
collagen.
26. (new) The method of claim 21 or 22 wherein the growth factor is epidermal
growth factor.
27. (new) The method of claim 21 or 22 wherein the growth factor is hepatocyte
growth factor.
28. (new) A population of hepatocytes and nonparenchymal cells, derived using a
method comprising:
co-culturing hepatocytes and nonparenchymal cells derived from
disaggregated liver tissue, in the presence of
(a) one or more growth factors that support the growth of
hepatocytes and
(b) a matrix wherein said matrix is a bead coated with at least one
biologically active molecule that promotes cell adhesion

under conditions sufficient to allow for the proliferation of said hepatocytes while retaining hepatic function of said hepatocytes.

29. (new) A population of hepatocytes and nonparenchymal cells, derived from a method comprising:

co-culturing established hepatic cell lines comprising hepatocytes and non-parenchymal cells, in the presence of

(a) one or more growth factors that support the growth of hepatocytes and

(b) a matrix wherein said matrix is a bead coated with at least one
biologically active molecule that promotes cell adhesion
under conditions sufficient to allow for the proliferation of said hepatocytes while
retaining hepatic function of said hepatocytes.—